



## Anti-HMG-CoA reductase autoantibodies

English text version in progress.

### Literature

Mohassel P, Mammen AL: Statin-associated autoimmune myopathy and anti-HMGCR autoantibodies. *Muscle Nerve* (2013); 48(4): 477 - 483 (PMID: [23519993](#)).

Stenzel W, Goebel HH, Aronica E: Review: immune-mediated necrotizing myopathies--a heterogeneous group of diseases with specific myopathological features. *Neuropathol Appl Neurobiol* (2012); 38(7): 632 - 646 (PMID: [22989019](#)).

Werner JL, Christopher-Stine L, Ghazarian SR, Pak KS, Kus JE, Daya NR, Lloyd TE, Mammen AL: Antibody levels correlate with creatine kinase levels and strength in anti-3-hydroxy-3-methylglutaryl-coenzyme A reductase-associated autoimmune myopathy. *Arthritis Rheum* (2012); 64(12): 4.087 - 4.093 (PMID: [22933019](#)).

Mammen AL, Pak K, Williams EK, Brisson D, Coresh J, Selvin E, Gaudet D: Rarity of anti-3-hydroxy-3-methylglutaryl-coenzyme A reductase antibodies in statin users, including those with self-limited musculoskeletal side effects. *Arthritis Care Res* (2012); 64(2): 269 - 272 (PMID: [21972203](#)).

Mammen AL, Chung T, Christopher-Stine L, Rosen P, Rosen A, Doering KR, Casciola-Rosen LA: Autoantibodies against 3-hydroxy-3-methylglutaryl-coenzyme A reductase in patients with statin-associated autoimmune myopathy. *Arthritis Rheum* (2011); 63(3): 713 - 721 (PMID: [21360500](#))

Christopher-Stine L, Casciola-Rosen LA, Hong G, Chung T, Corse AM, Mammen AL: A novel autoantibody recognizing 200-kd and 100-kd proteins is associated with an immune-mediated necrotizing myopathy. *Arthritis Rheum* (2010); 62(9): 2.757 - 2.766 (PMID: [20496415](#)).