



Anti-NXP-2 (MJ) autoantibodies

English text version in progress.

Literature

Ceribelli A, Fredi M, Taraborelli M, Cavazzana I, Franceschini F, Quinzanini M, Tincani A, Ross SJ, Chan JY, Pauley BA, Chan EK, Satoh M: Anti-MJ/NXP-2 autoantibody specificity in a cohort of adult Italian patients with polymyositis/dermatomyositis. *Arthritis Res Ther* (2012); 14(2): R97 (PMID: [22546500](#)).

Espada G, Maldonado Cocco JA, Fertig N, Oddis CV: Clinical and serologic characterization of an Argentine pediatric myositis cohort: identification of a novel autoantibody (anti-MJ) to a 142-kDa protein. *J Rheumatol* (2009); 36(11): 2.547 - 2.551 (PMID: [19833756](#)).

Fiorentino DF, Chung LS, Christopher-Stine L, Zaba L, Li S, Mammen AL, Rosen A, Casciola-Rosen L: Most patients with cancer-associated dermatomyositis have antibodies to nuclear matrix protein NXP-2 or transcription intermediary factor 1-gamma. *Arthritis Rheum* (2013); (PMID: [24037894](#)).

Gunawardena H, Wedderburn LR, Chinoy H, Betteridge ZE, North J, Ollier WE, Cooper RG, Oddis CV, Ramanan AV, Davidson JE, McHugh NJ: Autoantibodies to a 140-kd protein in juvenile dermatomyositis are associated with calcinosis. *Arthritis Rheum* (2009); 60(6): 1.807 - 1.814 (PMID: [19479859](#)).

Ichimura Y, Matsushita T, Hamaguchi Y, Kaji K, Hasegawa M, Tanino Y, Inokoshi Y, Kawai K, Kanekura T, Habuchi M, Igarashi A, Sogame R, Hashimoto T, Koga T, Nishino A, Ishiguro N, Sugimoto N, Aoki R, Ando N, Abe T, Kanda T, Kuwana M, Takehara K, Fujimoto M: Anti-NXP2 autoantibodies in adult patients with idiopathic inflammatory myopathies: possible association with malignancy. *Ann Rheum Dis* (2012) 71(5): 710 - 713 (PMID: [22258483](#)).

Ishikawa A, Muro Y, Sugiura K, Akiyama M: Development of an ELISA for detection of autoantibodies to nuclear matrix protein 2. *Rheumatology, Oxford* (2012); 51(7): 1.181 - 1.187 (PMID: [22427409](#)).

Oddis CV, Fertig N, Goel A, Espada G, Confalone Gregorian M, Maldonado Cocco JA: Clinical and serological characterization of the anti-MJ antibody in childhood myositis. *Arthritis Rheum* (1997); 40: S139.

Targoff IN, Trieu EP, Levy-Neto M, Fertig N, Oddis CV. Sera with autoantibodies to the MJ antigen react with NXP2: *Arthritis Rheum* (2007); 56: S787.