



Autoanticorpi anti-miosina

Prevalenza Cardiomiopatie dilatative, Miastenia gravis (fino a 32 %; Williams et al. 1986, 1987; Ohta et al. 1990; Takaya et al. 1992)

vedi

- ▶ [Autoanticorpi nelle malattie della giunzione neuromuscolare](#)
- ▶ [Autoanticorpi nelle malattie cardiovascolari](#)

Tabella 1 Malattie associate a anti-miosina

Malattie		Autori
Mistenia gravis	Prevalenza 23 %	Takaya et al. 1992 Ohta et al. 1990 Williams et al. 1987 Williams und Lennon 1986
Elenco di altre malattie associate in allestimento.		

Metodi di ricerca Il dosaggio degli anticorpi nel siero o plasma può essere ricercato fra l'altro con ELISA.

Bibliografia

Ohta M, Ohta K, Itoh N, Kurobe M, Hayashi K, Nishitani H: Anti-skeletal muscle antibodies in the sera from myasthenic patients with thymoma: identification of anti-myosin, actomyosin, actin, and alpha-actinin antibodies by a solid-phase radioimmunoassay and a western blotting analysis. Clin Chim Acta (1990); 187(3): 255 - 264 (PMID: [2323065](#)).

Takaya M, Kawahara S, Namba T, Grob D: Antibodies against myofibrillar proteins in myasthenia gravis patients. Tokai J Exp Clin Med (1992); 17(1): 35 - 39 (PMID: [1523691](#)).

Williams CL, Lennon VA, Momoi MY, Howard FM Jr. Serum antibodies and monoclonal antibodies secreted by thymic B-cell clones from patients with myasthenia gravis define striational antigens. Ann N Y Acad Sci. 1987;505:168-79. PubMed PMID: [3500666](#).

Williams CL, Lennon VA. Thymic B lymphocyte clones from patients with myasthenia gravis secrete monoclonal striational autoantibodies reacting with myosin, alpha actinin, or actin. J Exp Med. 1986 Oct 1;164(4):1043-59. PubMed PMID: [3020150](#); PubMed Central PMCID: PMC2188418.