



## Autoanticorpi anti-PIT-1

### Bibliografia

Bando H, Iguchi G, Yamamoto M, Hidaka-Takeno R, Takahashi Y: Anti-PIT-1 antibody syndrome; a novel clinical entity leading to hypopituitarism. *Pediatr Endocrinol Rev* (2015); 12(3): 290 - 296 (PMID: [25962206](#)).

Bando H, Iguchi G, Fukuoka H, Yamamoto M, Hidaka-Takeno R, Okimura Y, Matsumoto R, Suda K, Nishizawa H, Takahashi M, Tojo K, Takahashi Y: Involvement of PIT-1-reactive cytotoxic T lymphocytes in anti-PIT-1 antibody syndrome. *J Clin Endocrinol Metab* (2014); 99(9): E1.744 - 1.749 (PMID: [24937538](#)).

Yamamoto M, Iguchi G, Bando H, Fukuoka H, Suda K, Takahashi M, Nishizawa H, Matsumoto R, Tojo K, Mokubo A, Ogata T, Takahashi Y: A missense single-nucleotide polymorphism in the sialic acid acetyltransferase (SIAE) gene is associated with anti-PIT-1 antibody syndrome. *Endocr J* (2014); 61(6): 641 - 644 (PMID: [24748456](#)).

Yamamoto M, Iguchi G, Takeno R, Okimura Y, Sano T, Takahashi M, Nishizawa H, Handayaningshi AE, Fukuoka H, Tobita M, Saitoh T, Tojo K, Mokubo A, Morinobu A, Iida K, Kaji H, Seino S, Chihara K, Takahashi Y: Adult combined GH, prolactin, and TSH deficiency associated with circulating PIT-1 antibody in humans. *J Clin Invest* (2011); 121(1): 113 - 119 (PMID: [21123951](#)).