




## Autoanticorpi anti-solfato-3-glucoronil paragloboside (SGPG)

<b>Acronimo</b>	SGPG
<b>Omologo</b>	<b>SGLPG</b> (Solfato-3-glucoronil-lattosaminil-paragloboside)
<b>vedi</b>	<u>Autoanticorpi nelle neuropatie centrali</u> <u>Autoanticorpi in neuropatie periferiche</u>



**Figura 1** Solfato-3-glucoronil paragloboside (SGPG). Segnalato in rosso il resto del solfato dell'acido glucuronico.

### Bibliografia

- Ariga T: The role of sulfoglucuronosyl glycosphingolipids in the pathogenesis of monoclonal IgM paraproteinemia and peripheral neuropathy. Proc Jpn Acad. Ser B, Phys Biol Sci. 2011;87(7):386-404. Review. PubMed PMID: [21785257](#).
- Ariga T, Kohriyama T, Freddo L, Latov N, Saito M, Kon K, Ando S, Suzuki M, Hemling ME, Rinehart KL Jr, et al: Characterization of sulfated glucuronic acid containing glycolipids reacting with IgM M-proteins in patients with neuropathy. J Biol Chem (1987); 262(2): 848 - 853 (PMID: [2433270](#)).
- Ariga T, Suetake K, Nakane M, Kubota M, Usuki S, Kawashima I, Yu RK: Glycosphingolipid antigens in neural tumor cell lines and anti-glycosphingolipid antibodies in sera of patients with neural tumors. Neurosignals (2008); 16(2-3): 226 - 234 (PMID: [18253060](#)).
- Baumann N, Harpin ML, Marie Y, Lemerle K, Chassande B, Bouche P, Meininger V, Yu RK, Léger JM: Antiglycolipid antibodies in motor neuropathies. Ann N Y Acad Sci (1998); 845:322 - 329 (PMID: [9668365](#)).
- Bollensen E, Steck AJ, Schachner M: Reactivity with the peripheral myelin glycoprotein P0 in serum from patients with monoclonal IgM gammopathy and polyneuropathy. Neurology (1988); 38(8): 1.266 - 1.270 (PMID: [2456490](#)).
- Braun PE, Frail DE, Latov N: Myelin-associated glycoprotein is the antigen for a monoclonal IgM in polyneuropathy. J Neurochem (1982); 39(5): 1.261 - 1.265 (PMID: [6181197](#)).
- Caudie C, Vial C, Petiot P, Bancel J, Lombard C, Gonnaud PM: Monoclonal IgM autoantibody activity vis-à-vis glycoconjugates of peripheral nerves: apropos of 112 cases. Ann Biol Clin, Paris (2001); 59(5): 567 - 577 (PMID: [11602387](#)).
- Caudie C, Kaygisiz F, Jaquet P, Petiot P, Gonnaud PM, Antoine JC, Vial C: Diagnostic value of autoantibodies to MAG by ELISA Bühlmann in 117 immune-mediated peripheral neuropathies associated with monoclonal IgM to SGPG/SGLPG. Ann Biol Clin, Paris (2006); 64(4): 353 - 359 (PMID: [16829480](#)).
- Chassande B, Léger JM, Younes-Chennoufi AB, Bengoufa D, Maisonobe T, Bouche P, Baumann N: Peripheral neuropathy associated with IgM monoclonal gammopathy: correlations between M-protein antibody activity and clinical/electrophysiological features in 40 cases. Muscle Nerve (1998); 21(1):55 - 62 (PMID: [9427224](#)).



## Autoanticorpi anti-solfato-3-glucoronil paragloboside (SGPG)

Donfrid M, Apostolski S, Suvajdžić N, Janković G, Cemerikić-Martinović V, Atkinson HD, Colovic M: Monocytoid B cell lymphoma associated with antibodies to myelin-associated glycoprotein and sulphated glucuronyl paragloboside. *Acta Haematol* (2001); 106(3): 130 - 132 (PMID: [11713380](#)).

Gono T, Matsuda M, Shimojima Y, Ishii W, Yamamoto K, Morita H, Hashimoto T, Susuki K, Yuki N, Ikeda S: Rituximab therapy in chronic inflammatory demyelinating polyradiculoneuropathy with anti-SGPG IgM antibody. *J Clin Neurosci* (2006); 13(6): 683 - 687 (PMID: [16814550](#)).

Hashiguchi S, Ogasawara N, Shirabe T, Yoshino H: Axonal motor neuropathy associated with anti-SGPG antibody. *Rinsho Shinkeigaku* (1997); 37(3): 224 - 227 (PMID: [9217421](#)).

Ilyas AA, Gu Y, Dalakas MC, Quarles RH, Bhatt S: Induction of experimental ataxic sensory neuronopathy in cats by immunization with purified SGPG. *J Neuroimmunol* (2008); 193(1-2): 87 - 93 (PMID: [18037501](#)).

Ilyas AA, Mithen FA, Dalakas MC, Wargo M, Chen ZW, Bielory L, Cook SD: Antibodies to sulfated glycolipids in Guillain-Barré syndrome. *J Neurol Sci* (1991); 105(1): 108 - 117 (PMID: [1795163](#)).

Kuijff ML, Eurelings M, Tio-Gillen AP, van Doorn PA, van den Berg LH, Hooijkaas H, Stork J, Notermans NC, Jacobs BC: Detection of anti-MAG antibodies in polyneuropathy associated with IgM monoclonal gammopathy. *Neurology* (2009); 73(9): 688 - 695 (PMID: [19720975](#)).

McGinnis S, Kohriyama T, Yu RK, Pesce MA, Latov N: Antibodies to sulfated glucuronic acid containing glycosphingolipids in neuropathy associated with anti-MAG antibodies and in normal subjects. *J Neuroimmunol* (1988); 17(2): 119 - 126 (PMID: [2447123](#)).

Motoyama R, Yamakawa K, Suzuki S, Kusunoki S, Tanaka M: Rapid improvement by rituximab treatment in a case of demyelinating polyneuropathy with anti-myelin-associated glycoprotein antibody]. *Rinsho Shinkeigaku* (2011);51(10): 761 - 764 (PMID: [22019868](#)).

Nakamura H, Endo M, Sugawara E, Kuwahara M, Kusunoki S, Tanaka F, Takahashi T: A case of IgM paraproteinemic neuropathy associated with anti-sulfated glucuronic paragloboside (SGPG) IgG antibody without anti-myelin-associated glycoprotein (MAG) activity. *Rinsho Shinkeigaku* (2013); 53(10): 799 - 802 (PMID: [24225562](#)).

Rowland LP, Sherman WL, Hays AP, Lange DJ, Latov N, Trojaborg W, Younger DS: Autopsy-proven amyotrophic lateral sclerosis, Waldenström's macroglobulinemia, and antibodies to sulfated glucuronic acid paragloboside. *Neurology* (1995); 45(4): 827 - 829 (PMID: [7723980](#)).

Tagawa Y, Yuki N, Hirata K: Anti-SGPG antibody in CIDP: nosological position of IgM anti-MAG/SGPG antibody-associated neuropathy. *Muscle Nerve* (2000); 23(6): 895 - 899 (PMID: [10842265](#)).

Trojaborg W, Hays AP, van den Berg L, Younger DS, Latov N: Motor conduction parameters in neuropathies associated with anti-MAG antibodies and other types of demyelinating and axonal neuropathies. *Muscle Nerve* (1995); 18(7): 730 - 735 (PMID: [7540258](#)).

Van den Berg L, Hays AP, Nobile-Orazio E, Kinsella LJ, Manfredini E, Corbo M, Rosoklija G, Younger DS, Lovelace RE, Trojaborg W, Lange DE, Goldstein S, Delfiner JS, Sadiq SA, Sherman WH, Latov N: Anti-MAG and anti-SGPG antibodies in neuropathy. *Muscle Nerve* (1996); 19(5): 637 - 643 (PMID: [8618562](#)).

Weiss MD, Dalakas MC, Lauter CJ, Willison HJ, Quarles RH: Variability in the binding of anti-MAG and anti-SGPG antibodies to target antigens in demyelinating neuropathy and IgM paraproteinemia. *J Neuroimmunol* (1999); 95(1-2): 174 - 184 (PMID: [10229128](#)).